

TITLE: Evaluation Of Spent Coffee Obtained From The Most Common Coffeemakers As A Source Of Hydrophilic Bioactive Compounds

AUTHORS: Jimena Bravo^a, Isabel Juárez^a, Carmen Monente^a, Bettina Caemmerer^b, Lothar W. Kroh^b, M. Paz De Peña^{a*} and Concepción Cid^a

^aDepartment of Nutrition, Food Science and Physiology, School of Pharmacy, University of Navarra, E-31008-Pamplona, Spain.

^bInstitut für Lebensmittelchemie und Lebensmitteltechnologie, Technische Universität Berlin, Gustav-Meyer-Allee 25, D-13355 Berlin, Germany.

Published in

Journal of Agricultural and Food Chemistry 60, 12565-12573 (2012)

*Corresponding author:

M. Paz De Peña

Department of Nutrition, Food Science and Physiology

School of Pharmacy, University of Navarra

C/ Irunlarrea 1

E-31080-Pamplona, Spain

Tel: +34 948 425600 (806580)

Fax: +34 948 425740

E-mail address: mpdepena@unav.es

ABSTRACT

The main hydrophilic antioxidant compounds (3-, 4- and 5-monocaffeoylquinic and 3,4-, 3,5- and 4,5-dicaffeoylquinic acids, caffeine and browned compounds, including melanoidins) and the antioxidant capacity (Folin-Ciocalteu, ABTS, DPPH, Fremy's salt and TEMPO) were evaluated in Arabica and Robusta spent coffee obtained from the preparation of coffee brews with the most common coffeemakers (filter, espresso, plunger and mocha). All spent coffee grounds, with the exception of those from mocha coffeemaker, had relevant amounts of total caffeoylquinic acids (6.22–13.24 mg/g of spent coffee), mainly dicaffeoyquinic acids (3.31-5.79 mg/g of spent coffee) which were 4-7 fold higher than in their respective coffee brews. Caffeine was ranged from 3.59 to 8.09 mg/g of spent coffee. The antioxidant capacity of the aqueous spent coffee extracts was 46.0-102.3% (filter), 59.2-85.6% (espresso), and less than 42% (plunger) in comparison to their respective coffee brews. This study allowed us to obtain spent coffee extracts with antioxidant properties that can be used as a good source of hydrophilic bioactive compounds.

KEYWORDS:

Coffee; Spent Coffee; By-products; Bioactive Compounds; Chlorogenic Acids; Antioxidant

Introduction

Millions of coffee cups are consumed every day around the world producing spent coffee grounds in tons in restaurants and cafeterias, but also at domestic levels. Traditionally these residues have been considered as wastes, or occasionally used as fertilizer. During last decades, at industrial level several applications have been proposed to add value to coffee by-products originated from the fruit to roasted coffee (coffee pulp, husk, and silverskin) and even soluble coffee (spent coffee).^{1, 2} For example, spent coffee from soluble coffee industry might be used for animal feed,³ biofuel, biodiesel or bioethanol production,^{4, 5} as adsorbent and activated carbon,^{6, 7} but also to extract antioxidants or other bioactive compounds.^{8, 9}

Although the production of soluble (or instant) coffee originates a high effective extraction of coffee bioactive compounds, such as caffeine, chlorogenic acids, etc., limited amounts of these compounds still remains in spent coffee grounds and might be extracted with solvents (ethanol, methanol, etc.).^{8, 10} Similarly, the spent coffee grounds originated by the preparation of a cup of coffee might also contain relevant amounts of bioactive compounds. However, a direct transfer of the results of industry soluble coffee extraction to coffee brew preparation is not possible because, despite the type of coffee, there is a clear influence of (i) different coffee brewing procedures, including different coffeemakers,^{11, 12} but also (ii) technological factors, such as water temperature and pressure, coffee/water ratio, grinding, etc. on the extraction of coffee compounds.¹³⁻¹⁶ Currently, to our best knowledge, only few studies about the presence of phenolics in spent coffee grounds from espresso coffee,^{17, 18} and the preparation of extracts with antioxidant capacity from spent coffee grounds obtained by filter coffee brewing have been found.¹⁹

On the other hand, the increasing demand for foodstuffs free of artificial additives or with nutritional or healthy added values, induces to food industry (and also to the pharmaceutical one) to find new sources of antioxidants and bioactive compounds. For that reason, the knowledge of the main bioactive compounds extracted from spent coffee grounds would give them a higher value. Therefore, the aim of this work was to quantitate the main bioactive compounds (caffeoylquinic acids, caffeine and browned compounds, including melanoidins) in spent coffee obtained from the preparation of coffee brews with the most common coffeemakers (filter, espresso, plunger and mocha), and to evaluate their antioxidant capacity. This study allows us to know whether the residues generated during coffee brewing procedure, produced in large amounts in cafeterias and restaurants, or at domestic levels, can be considered as a source of natural antioxidants.

Materials and methods

Chemicals and reagents

The methanol used (spectrophotometric and HPLC grade) and Folin-Ciocalteu reagent were from Panreac (Barcelona, Spain). ABTS (2,2'-Azino-bi(3-ethylbenzo-thiazonile-6-sulfonic acid) diammonium salt), potassium persulfate, DPPH• (2,2-Diphenyl-1-picrylhydrazyl), Trolox (6-hydroxy-2,5,7,8-tetramethyl-chroman-2-carboxylic acid), dipotassium hydrogen phosphate, potassium dihydrogen phosphate, sodium chloride, Fremy's salt (potassium nitrosodisulfonate) and TEMPO (2,2,6,6-tetramethyl-1-piperidin-1-oxyl) were purchased from Sigma-Aldrich (Steinheim, Germany). Gallic acid was from Fluka (Madrid, Spain). Pure reference standards of 5-caffeoylquinic acid (5-CQA) and caffeine were obtained from Sigma-Aldrich (Steinheim, Germany) and pure reference standards of 3,4-, 3,5- and 4,5-dicafeoylquinic acids were purchased

from Phytolab (Vestenbergsgreuth, Germany). A mixture of 3-CQA, 4-CQA, and 5-CQA was prepared from 5-CQA using the isomerization method of Trugo & Macrae,²⁰ also described in Farah et al.²¹

Spent Coffee Preparation

Roasted coffee (without defective beans) from Guatemala (*Coffea arabica*, named Arabica, 3.03 % water content, $L^* = 25.40 \pm 0.69$, roasted at 219 °C for 905 s) and Vietnam (*Coffea canephora* var. robusta, named Robusta, 1.59 % water content, $L^* = 24.92 \pm 0.01$, roasted at 228 °C for 859 s) was provided by a local factory. Coffee beans were ground for 20 s using a grinder (model Moulinex super junior “s”, Paris, France). The L^* value was analyzed by means of a tristimulus colorimeter (Chromameter-2 CR-200, Minolta, Osaka, Japan) using the D65 illuminant and CIE 1931 standard observer. The instrument was standardized against a white tile before sample measurements. Ground roasted coffee was spread out in a 1 cm Petri plate, and the L^* value was measured in triplicate on the CIELab scale. Water content was measured by weight loss after drying for 2 h at 102 ± 3 °C in an oven JP SELECTA (Barcelona, Spain).

Four coffee brewing procedures were selected to obtain the spent coffee: filter coffeemaker, espresso coffeemaker, plunger coffeemaker and mocha coffeemaker. The coffee brews were prepared as described by López-Galilea et al.¹¹ with some modifications. Filter coffee brew was prepared from 24 g of ground roasted coffee for a volume of 400 mL of water, using a filter coffeemaker (model AVANTIS 70 Inox, Ufesa, Spain). Extraction took approximately 6 min at 90 °C. Espresso coffee brew was prepared from 7 g of ground roasted coffee for a volume of 40 mL using an espresso coffeemaker (model Saeco Aroma, Italy). The water pump pressure was 15 bar. Extraction took approximately 24 s at 90 °C. Plunger coffee brew was prepared from 40 g of ground roasted coffee, which was extracted adding 500 mL of water at 98 °C in a

plunger coffeemaker (model Bistro Nouveau Coffee Maker, Bodum, 1L capacity). The water and the coffee powder were kept in contact for 5 min before the plunger was slowly pushed down. Mocha coffee brew was prepared from 36 g of ground roasted coffee for a volume of 450 mL, using a mocha coffeemaker (model Bra, Spain). The heating temperature and extraction time were approximately 10 min at 93 °C.

Ground roasted coffee after brewing procedures, called spent coffee, were dried to a constant weight for 2 hours at 102 ±3 °C in an oven JP SELECTA (Barcelona, Spain).

Spent Coffee Extracts Preparation

Spent Coffee extracts were prepared according to the method described by Bravo et al.¹⁹ Briefly, first, dried spent coffee was defatted with Petroleum Ether (1:11, w/v) for 3 h at 60 °C in a Soxhlet extraction system (Extraction Unit B-811 Standard BÜCHI, Flawil, Switzerland). Then, 24 g of spent coffee were extracted with a volume of 400 mL of water using a filter coffeemaker (model AVANTIS 70 Inox, Ufesa, Spain). Extraction took approximately 6 min at 90 °C. Aqueous spent coffee extracts were lyophilized using a CRYODOS Telstar (Terrassa, Spain).

Caffeoylquinic acids and caffeine

Extract preparation and cleanup were carried out according to Bicchi et al.²² The compounds were analyzed by HPLC following the method described by Farah et al.,²¹ with some modifications. HPLC analysis was achieved with an analytical HPLC unit model 1100 (Agilent Technologies, Palo Alto, CA, USA) equipped with a binary pump and an automated sample injector. A reversed-phase Hypersil-ODS (5 µm particle size, 250 x 4.6 mm) column was used at 25 °C. The sample injection volume was 100 µL. The chromatographic separation was performed using a gradient of methanol (solvent A) and Milli-Q water acidulated with phosphoric acid (pH 3.0, solvent B) at a constant flow of 0.8 mL/min starting with 20% solvent A. Then solvent A was increased to 50%

within 15 min to be maintained at 50% for 9 min and, finally, to return to initial conditions (20% solvent A) in 3 min. Detection was accomplished with a diode-array detector, and chromatograms were recorded at 325 nm for caffeoylquinic acids and 276 nm for caffeine. Identification of caffeoylquinic acids and caffeine was performed by comparing the retention time and the UV/VIS spectra with those of their reference compounds. Quantitation of 5-caffeoylquinic (5-CQA) and caffeine was made by comparing the peak areas with those of the standards.

The method was validated obtaining a linear relationship between the concentrations of each compound and the respective UV absorbance ($r=0.999$). The recovery values were $101.3\pm 1.3\%$ for 5-CQA and $106.3\pm 2.4\%$ for caffeine. Results for repeatability showed good precision of the method with coefficient of variation values below 5%. A narrow dispersion of values was also observed for intermediate precision, with coefficients of variation between 0.11% and 4.96%. The detection limits were 1.18 $\mu\text{g/mL}$ and 0.83 $\mu\text{g/mL}$ for 5-CQA and caffeine, respectively. The quantitation limits were 3.95 $\mu\text{g/mL}$ for 5-CQA and 2.76 $\mu\text{g/mL}$ for caffeine.

Quantitation of the other caffeoylquinic acids was performed using the area of 5-CQA standard combined with molar extinction coefficients of the respective caffeoylquinic acid as reported by Trugo et al.²⁰ and Farah et al.,²¹ using the following equation:

$$C = \frac{RF \times \varepsilon_1 \times M_{r2} \times A}{\varepsilon_2 \times M_{r1}}$$

where C is the concentration of the isomer in milligrams per milliliter; RF is the response factor of the 5-CQA standard (concentration in milligrams per milliliter per unit area); ε_1 is the molar extinction coefficient of 5-CQA; ε_2 is the molar extinction coefficient of the isomer in question; M_{r1} is the relative molecular mass of 5-CQA; M_{r2} is the relative molecular mass of the isomer in question; and A is the area of the peak corresponding to the isomer in question. Molar extinction coefficients ($\times 10^4$) were as

follows: at λ_{\max} 330 nm, 5-CQA = 1.95, 4-CQA = 1.80, 3-CQA = 1.84, 3,4-diCQA = 3.18, 3,5-diCQA = 3.16 and 4,5-diCQA = 3.32.

Results were expressed as milligrams of each compound per gram of spent coffee dry matter (spent coffee extracts) or per gram of coffee (coffee brews).

Browning Index (Abs 420 nm)

Fifty microliters of the sample (spent coffee extract or coffee brew) was diluted up to 2 mL with demineralized water. Browning index was measured by reading the absorbance of samples at 420 nm, after exactly 2 minutes in a 3 mL capacity cuvette (1 cm length) with a Lambda 25 UV-vis spectrophotometer (Perkin-Elmer Instruments, Madrid, Spain) connected to a thermostatically controlled chamber (25 °C) and equipped with UV WinLab software (Perkin Elmer). This index measures the browned compounds developed during caramelization and Maillard Reactions, including melanoidins.²³

Antioxidant Capacity by Folin-Ciocalteu (FC) assay

The Folin-Ciocalteu reducing capacity of coffee was performed according to the Singleton's method.²⁴ Spent coffee extracts and coffee brews were diluted 3:10 and 1:10 respectively in demineralized water prior to analysis. A volume of 500 μ L of Folin-Ciocalteu reagent (Panreac) was added to a mixture of 100 μ L of the extract sample and 7.9 mL of demineralized water. After a 2 min delay, 1.5 mL of a 7.5% sodium carbonate solution was added. Next, the sample was incubated in darkness at room temperature for 90 min. The absorbance of the sample was measured at 765 nm in a Lambda 25 UV/VIS spectrophotometer (Perkin Elmer Instruments, Madrid, Spain). Gallic acid (GA) was used as reference, and the results were expressed as milligrams of GA per gram of spent coffee dry matter (mg GA/g spent coffee dm) or per gram of coffee (mg GA/g coffee).

Antioxidant Capacity by ABTS Assay

The ABTS antioxidant capacity was performed according to the method of Re et al.²⁵ with some modifications. The radicals $ABTS^{\cdot+}$ were generated by the addition of 0.36 mM potassium persulfate to a 0.9 mM ABTS solution prepared in phosphate buffered saline (PBS) (pH 7.4), and the $ABTS^{\cdot+}$ solution was stored in darkness for 12 h. The $ABTS^{\cdot+}$ solution was adjusted with PBS to an absorbance of 0.700 (\pm 0.020) at 734 nm in a 3 mL capacity cuvette (1 cm length) at 25 °C (Lambda 25 UV-VIS spectrophotometer, Perkin-Elmer Instruments, Madrid, Spain). Spent coffee extracts and coffee brews were diluted 3:100 and 15:1000 respectively in demineralized water prior to analysis. Samples (50 μ L) were added to 2 mL of $ABTS^{\cdot+}$ solution. The absorbance was measured spectrophotometrically at 734 nm after exactly 18 min. Calibration was performed with Trolox solution (a water-soluble vitamin E analogue), and the antioxidant capacity was expressed as micromoles of Trolox per gram of spent coffee dry matter (μ mol Trolox /g spent coffee dm) or per gram of coffee (μ mol Trolox /g coffee).

Antioxidant Capacity by DPPH Assay

The antioxidant capacity was also measured using 2, 2-diphenyl-1-picrylhydrazyl DPPH \bullet decolorization assay (Brand-Williams et al.²⁶ with modifications). A 6.1×10^{-5} mol/L DPPH \bullet methanolic solution was prepared immediately before use. The DPPH \bullet solution was adjusted with methanol to an absorbance of 0.700 (\pm 0.020) at 515 nm in a 3 mL capacity cuvette (1 cm length) at 25 °C (Lambda 25 UV-VIS spectrophotometer, Perkin-Elmer Instruments, Madrid, Spain). Spent coffee extracts and coffee brews were diluted 3:100 and 15:1000 respectively in demineralized water prior to analysis. Samples (50 μ L) were added to 1.95 mL of the DPPH \bullet solution. After mixing, the absorbance was measured at 515 nm after exactly 18 min. Calibration was performed with Trolox solution (a water-soluble vitamin E analogue). The antioxidant capacity

was expressed as micromoles of Trolox per gram of spent coffee dry matter (μmol Trolox /g spent coffee dm) or per gram of coffee (μmol Trolox /g coffee).

Antioxidant Capacity by Electro Spin Resonance (ESR) Spectroscopy

The ESR spectroscopy measurements were performed with Fremy's salt and TEMPO as stabilized radicals with the same procedure as described by Roesch et al.²⁷ and modified by Cämmerer & Kroh.²⁸ For the investigation with Fremy's salt, 100 μL of every spent coffee extract diluted 100-fold with demineralized water were allowed to react with an equal volume of an aqueous 1 mM Fremy's salt solution prepared in 50 mM phosphate buffer (pH 7.4). ESR spectra were recorded every 35 s for 30 min. For the investigation with TEMPO, aliquots of 200 μL of spent coffee extract were allowed to react with 100 μL of 1 mM TEMPO solution. ESR spectra were obtained after 30 min, by which time the reaction was complete. Microwave power was set at 10 dB. Modulation amplitude, center field, and sweep width were set at 1.5, 3397, and 71 G, respectively. Both Fremy's salt and TEMPO antioxidant activity were calculated as Trolox equivalents and expressed as micromoles of Trolox per gram of spent coffee dry matter (μmol Trolox /g spent coffee dm).

Statistical Analysis

Each parameter was analyzed in triplicate. Results are shown as means \pm standard deviations. One-way Analysis of Variance (ANOVA) was applied for each parameter; the source of variation was the different brewing procedure for obtaining spent coffee. A Tukey test was applied as a test *a posteriori* with a level of significance of 95 %. All statistical analyses were performed using the SPSS v.15.0 software package.

Results and discussion

Caffeoylquinic acids

Caffeoylquinic acids are the most abundant phenolic compounds in coffee. These chlorogenic acids are water soluble esters formed between quinic acid and one or two moieties of caffeic acid, a *trans*-cinnamic acid. Monocaffeoylquinic acids (3-CQA, 4-CQA, 5-CQA) and dicaffeoylquinic acids (3,4-diCQA, 3,5-diCQA, 4,5-diCQA) were identified and quantified in spent coffee obtained from different coffeemakers (filter, espresso, plunger and mocha) and in their respective coffee brews. The results are shown in **Table 1**.

All spent coffee grounds, with the exception of those from mocha coffeemaker, had relevant amounts of total caffeoylquinic acids ranging from 11.05 (espresso) to 13.24 (filter) mg per g of Arabica spent coffee, and from 6.22 (filter) to 7.49 (espresso) mg per g of Robusta spent coffee. These amounts correspond to 56.6-86.6% of those caffeoylquinic acids obtained for the coffee brews. In espresso coffee residues, Cruz et al.¹⁷ reported similar amounts of 5-CQA. However, in the present study the total CQAs and diCQAs range values reach higher results than total CGA in Cruz et al. study.¹⁷ This discrepancy could be probably due to the different type of coffee, Arabica or Robusta in the present study or Arabica/Robusta blends in Cruz et al. work,¹⁷ but also to other technological factors, such as roasting degree, extraction methodology, etc. Despite these results, to our knowledge, the present study is the first one to report the quantity of individual mono and dicaffeoylquinic acids present in spent coffee obtained from coffeemakers different to espresso ones.

Coffee brews prepared with mocha coffeemaker showed the highest content in both monocaffeoylquinic (23.68 and 13.71 mg per g for Arabica and Robusta, respectively) and dicaffeoylquinic (1.94 and 1.58 mg per g for Arabica and Robusta, respectively) acids, suggesting a practically complete extraction of caffeoylquinic acids, being negligible the amount present in the spent coffee grounds. In a previous work, we also

observed that a second successive aqueous extraction from the spent coffee grounds only contributed to a few amount of phenolic compounds (less than 15% of those obtained in the first extraction).¹⁹ Therefore, it could be considered that the total content of caffeoylquinic acids present in coffee has been practically extracted with the preparation of the coffee brew and the first aqueous extraction of spent coffee. This was also corroborated by the fact that the sum of the total caffeoylquinic acids of each coffee brew and its respective spent coffee was quite similar among different coffeemakers ranged from 15.29 (mocha) to 16.29 (plunger) mg per g for Arabica and from 25.62 (mocha) to 30.77 (plunger) mg per g for Robusta coffee. These results are also in agreement with the total chlorogenic acids amounts reported for roasted coffees from different origins and varieties.^{29, 30} Although higher amounts of caffeoylquinic acids in Robusta coffees than in Arabica ones have been extensively reported,²¹ other authors have found less amounts of CQAs in Robusta coffees.^{31, 32} This could be because both, the origin of coffee and the higher loss of chlorogenic acids in Robusta coffee during roasting process.^{33, 34}

In spent coffee, 5-CQA and 3,4-diCQA were the most abundant caffeoylquinic acids, whereas 3,5-diCQA and 3-CQA were the least ones. Moreover, the content of total diCQAs was quite similar and even higher to total monoCQAs. This is due to 3 to 6 fold, and even higher for filter coffeemaker, amounts of total diCQAs in spent coffee in comparison to their respective coffee brews. In contrast, in coffee brews the monoCQAs content was 12 fold or even higher than diCQAs, being 5-CQA the most and 3,4-diCQA the least abundant. Therefore, the monoCQAs are mainly extracted during coffee brewing whereas the diCQAs need further conditions to be extracted. Other authors also observed that diCQAs or dicaffeoylquinides were extracted rather slowly from coffee in comparison to monoCQAs.^{32, 35} The esterification of an additional caffeic acid moiety in

diCQAs increases the number of hydroxyl groups that might be bound to the amide carbonyls of the peptide bond in melanoidins.^{36, 37} Also, other authors³⁸ corroborated this hypothesis that the chlorogenic acids are incorporated into melanoidins via hydroxycinnamic acids (such as caffeic acid) moieties, mainly through nonester linkages upon coffee roasting. Therefore, the release of diCQAs bound to melanoidins is rather difficult to that of monoCQAs. However, as proposed by Ludwig et al.,³² the use of an extraction system with turbulences during few minutes facilitates the contact of grounds and water, a polar solvent, and favors the caffeoylquinic acids extraction, also diCQAs. Taking into account all these considerations, the application of the appropriate conditions allowed us to obtain spent coffee extracts that can be considered as a good source of caffeoylquinic acids, mainly diCQAs.

Caffeine

The presence of caffeine in spent coffee grounds has been proposed as one of the main problems to be used in agriculture because its toxicity.³⁹ Although some few studies reported the content of caffeine in spent coffee,^{8, 10, 17} to our knowledge, the present study is the first one to report the amount of caffeine in spent coffee grounds from coffeemakers different to espresso ones. The results of the amount of caffeine in spent coffee and in their respective coffee brews are shown in **Table 2**.

All spent coffee grounds, with the exception of those from mocha coffeemaker, had concentrations of caffeine ranging from 3.59 (espresso) to 5.20 (filter) mg per g of Arabica spent coffee and from 5.73 (plunger) to 8.09 (espresso) mg per g of Robusta spent coffee. Quite similar amounts of caffeine were reported by Cruz et al.¹⁷ for espresso coffee residues obtained from Arabica/Robusta blends. The caffeine amounts in spent coffee were 2 to 3 fold lower to those obtained for their respective coffee brews. Taking into account that a second successive aqueous extraction of spent coffee

grounds extracted less than 3% of the caffeine concentration found in the first extraction, the total content of caffeine present in roasted coffee has been practically extracted with the preparation of the coffee brew and the first aqueous extraction of spent coffee. In fact, the sum of caffeine in coffee brews and spent coffee ranged from 13.66 to 15.64 mg per g for Arabica coffee and from 21.37 to 29.61 mg per g for Robusta coffee. These results are in agreement with those of the literature which extensively reports that robusta coffees are richer in caffeine than Arabica ones.⁴⁰

Similar to caffeoylquinic acids, mocha coffeemaker extracted the majority of the caffeine present in coffee, being the mocha coffee brews the richest in caffeine and leaving not detectable caffeine amounts in the spent coffee grounds. Consequently, spent coffee grounds from mocha coffeemaker and spent coffee grounds from the other coffeemakers after one extraction with water can be considered as caffeine-free coffee by-products.

Browned compounds

Browned compounds, mainly melanoidins, are originated by Maillard reactions during coffee roasting process. They contribute to the antioxidant and other functional properties of coffee by themselves and because of the presence of phenolics in their skeleton.^{41, 42} Spent coffee extracts obtained from filter coffeemaker showed the highest browning index (**Table 2**; 0.165 and 0.145 for Arabica and Robusta coffees, respectively). These results were similar to those of the filter coffee brews. In contrast, browned compounds in spent coffee extracts from espresso and plunger coffeemakers were 3 to 5 fold lower than in their respective coffee brews. Similar to the other bioactive compounds (caffeoylquinic acids and caffeine), aqueous soluble browned compounds were mainly extracted in coffee brews when mocha coffeemaker was used, releasing only traces in spent coffee. Yen et al.¹⁰ also reported a lower browning index in

aqueous extracts from soluble spent coffee grounds in comparison to that from roasted coffee. Therefore, although melanoidins and other browned compounds may act as antioxidants, because of their lower amounts in spent coffee, their contribution to the antioxidant capacity could be rather limited in comparison to that in coffee brews, where they reach up to an 80%.⁴³

Antioxidant capacity of spent coffee

The presence of antioxidant compounds, such as chlorogenic acids, caffeine and Maillard Reaction Products, have been extensively associated with the health benefits of coffee, such as cardiovascular and neurodegenerative diseases prevention, anti-carcinogenic and antimutagenic activity.^{44, 45} Beside these healthy properties, antioxidant capacity may contribute to the preservation of coffee against oxidative damage throughout the storage time. Antioxidant activity has been also reported by other authors in extracts of spent coffee grounds from the soluble coffee production,^{2, 8, 46} from espresso coffee^{17, 18} and from filter coffee in a previous work of our group.¹⁹ However, to our knowledge, the antioxidant capacity of spent coffee grounds obtained from other different coffeemakers (plunger and mocha) using an Arabica and a Robusta coffee has not been reported previously. For those reasons, the antioxidant capacity was evaluated using three different colorimetric assays (Folin Ciocalteu, ABTS and DPPH) in spent coffee obtained from different coffeemakers (filter, espresso, plunger and mocha) and in their respective coffee brews. The results of the antioxidant capacity are shown in **Table 3**.

As it was expected because of the presence of phenolic and nonphenolic antioxidants, all spent coffee, except those from mocha coffeemaker, showed relevant antioxidant capacity. Arabica spent coffee obtained from filter coffeemaker showed the highest values in all assays, followed by Robusta spent coffees from espresso and filter

coffeemakers, and Arabica spent coffee from espresso coffeemaker, in order. The results of the Folin-Ciocalteu assay of spent coffee from espresso coffeemaker (15.79 and 19.12 mg GA per gram of spent coffee for Arabica and Robusta coffees, respectively) were similar to those found by other authors.¹⁸ On the other hand, spent coffee obtained from plunger coffeemaker showed less antioxidant capacity, and no significant differences between Arabica and Robusta coffees were found.

These antioxidant capacity results seem to be due to a balance between caffeoylquinic acids, more abundant in Arabica spent coffee, and caffeine and browned compounds, mainly present in Robusta one. In fact, caffeine was in higher amounts in Robusta spent coffee than in Arabica, being Robusta spent coffee from espresso coffeemaker that with the highest content (8.09 mg per gram of spent coffee). Other authors also found high correlations between antioxidant capacity of coffee brews and caffeine, suggesting that caffeine might be a good contributor to the antioxidant capacity or reducing power of coffee brews^{11, 31, 32}, whereas other authors⁴³ find it for the melanoidins content.

The comparison between the antioxidant capacity of the spent coffee with their respective coffee brews (expressed as 100%) is shown in **Figure 1**. The antioxidant capacity of spent coffee obtained from filter coffeemaker was quite similar (88.9-102.3%) to that of the coffee brew for Arabica coffee, whereas for Robusta coffee was only 46.0-57.8%. However, for spent coffee grounds obtained with espresso coffeemaker, the antioxidant capacity was 59.2-68.7% and 64.0-85.6% of the Robusta and Arabica coffee brews, respectively. These results might be partially explained by the higher browning index in both espresso coffee brews (0.315 and 0.418), and in less extent in Robusta filter coffee brew (0.197) that reduce the amount of melanoidins and other browned compounds to be extracted from spent coffee. Moreover, it should be also taken into account that other minor phenolic (hydroxycinnamic acids,

feruloylquinic acids and other chlorogenic acids) and nonphenolic (volatiles, etc.) compounds might contribute to the antioxidant capacity. Thus, to go deeper into the knowledge of the contribution of phenolic and nonphenolic compounds to the antioxidant capacity of spent coffee, Electro Spin Resonance (ESR) spectroscopy was used to measure the antioxidant capacity of spent coffee obtained from filter and espresso coffeemakers.

Mainly phenolic compounds can be detected when Fremy's salt is used as the stabilized radical, whereas TEMPO radical is mainly scavenged by Maillard reaction products (MRP), such as melanoidins.^{12, 28} As it is shown in **Figure 2**, ESR antioxidant capacity of Arabica spent coffee from espresso coffeemaker was lower than that from filter one, whereas for Robusta spent coffee these results were reversed. The behaviour of the results of antioxidant capacity measured by Fremy's salt technique (**Figure 2a**) was similar to those obtained by the colorimetric assays, but rather different to that of the amounts of caffeoylquinic acids in spent coffee. This may be explained because, although caffeoylquinic acids are the most abundant phenolics in coffee, other minor hydroxycinnamic acids, such as caffeic acid, ferulic acid and its derivatives by esterification with quinic acid, might also contribute to the Fremy's salt antioxidant capacity of spent coffee. Moreover, although Fremy's salt is mainly scavenged by chlorogenic acids, other antioxidants, including melanoidins, can scavenge this stabilized radical in a nonspecific manner and in a very low proportion.^{12, 47} On the other hand, TEMPO scavenging capacity (**Figure 2b**) was at least 10 fold lower than Fremy's salt, showing that the contribution of roasting-induced antioxidants is rather limited in spent coffee. Similar behaviour was also observed in coffee brews by other authors.^{12, 32, 47} For that reason, all TEMPO results should be considered as very low, including the significantly higher TEMPO antioxidant capacity of the aqueous extracts

obtained from Arabica spent coffee grounds from filter coffeemaker. This higher result might be explained by the slightly higher browning index (Abs 420nm 0.165), but also because melanoidins have different antioxidant capacities, depending on the structure, their composition and the phenolics bound in their skeleton.^{12, 47-49}

In conclusion, spent coffee obtained from the most common coffeemakers used at domestic and cafeterias level (filter and espresso), and in less proportion from plunger ones, could be considered as a good potential source of hydrophilic bioactive compounds. These compounds are easily extracted with water to obtain extracts with antioxidant properties that might be used in food and pharmaceutical industries to increase food nutritional values, product stability or to develop new healthy products. In contrast, spent coffee grounds from mocha coffeemaker only have traces of hydrophilic bioactive compounds because this coffeemaker is very effective to extract them in coffee brews. Moreover, these spent coffee grounds and the spent coffee grounds from the other coffeemakers after one extraction with water, might be subsequently used for agriculture because they are practically caffeine-free coffee by-products. Therefore, this study allows us to know that it is possible to consider spent coffee as an added-value by-product because it can be used as a good source of bioactive compounds and after that in agriculture. However, further research should be needed to identify and quantitate the minor antioxidant compounds present in spent coffee and to demonstrate their functional or healthy properties.

Acknowledgements

This research was funded by the Spanish Ministry of Economy and Competitiveness (AGL2009-12052). We thank the Spanish Ministry of Education (PR2009-0324), the Association of Friends of the University of Navarra and the Department of Industry of

the Government of Navarre for the grants given to M.P.P., J.B., C.M. and I.J., respectively. We thank the Unión Tostadora S.A. for providing the coffee.

This study was partially presented at the 1st International Congress on Cocoa, Coffee and Tea, September 2011, Novara (Italy), and at the 5th International Conference on Polyphenols and Health, October 2011, Sitges (Spain).

References

1. Esquivel, P.; Jiménez, V.M. Functional properties of coffee and coffee by-products. *Food Res. Int.* **2012**, *46*, 488-495.
2. Murthy, P.; Madhava Naidu, M. Recovery of Phenolic Antioxidants and Functional Compounds from Coffee Industry By-Products. *Food Bioprocess Technol.* **2012**, *5*, 897-903.
3. Givens, D.I.; Barber, W.P. In vivo evaluation of spent coffee grounds as a ruminant feed. *Agric. Wastes* **1986**, *18*, 69-72.
4. Silva, M.; Nebra, S.; Machado Silva, M.; Sanchez, C. The use of biomass residues in the Brazilian soluble coffee industry. *Biomass Bioenergy* **1998**, *14*, 457-467.
5. Kondamudi, N.; Mohapatra, S.K.; Misra, M. Spent Coffee Grounds as a Versatile Source of Green Energy. *J. Agric. Food Chem.* **2008**, *56*, 11757-11760.
6. Namane, A.; Mekarzia, A.; Benrachedi, K.; Belhaneche-Bensemra, N.; Hellal, A. Determination of the adsorption capacity of activated carbon made from coffee grounds by chemical activation with ZnCl₂ and H₃PO₄. *J. Hazard. Mater.* **2005**, *119*, 189-194.
7. Franca, A.S.; Oliveira, L.S.; Ferreira, M.E. Kinetics and equilibrium studies of methylene blue adsorption by spent coffee grounds. *Desalination* **2009**, *249*, 267-272.
8. Ramalakshmi, K.; Rao, L.J.M.; Takano-Ishikawa, Y.; Goto, M. Bioactivities of low-grade green coffee and spent coffee in different in vitro model systems. *Food Chem.* **2009**, *115*, 79-85.
9. Mussatto, S.I.; Ballesteros, L.F.; Martins, S.; Teixeira, J.A. Extraction of antioxidant phenolic compounds from spent coffee grounds. *Sep. Purif. Technol.* **2011**, *83*, 173-179.
10. Yen, W. J.; Wang, B.; Chang, L.; Duh, P. Antioxidant Properties of Roasted Coffee Residues. *J. Agric. Food Chem.* **2005**, *53*, 2658-2663.
11. López-Galilea, I.; de Peña, M.P.; Cid, C. Correlation of selected constituents with the total antioxidant capacity of coffee beverages: influence of the brewing procedure. *J. Agric. Food Chem.* **2007**, *55*, 6110-6117.
12. Pérez-Martínez, M.; Caemmerer, B.; De Peña, M.; Cid, C.; Kroh, L. Influence of Brewing Method and Acidity Regulators on the Antioxidant Capacity of Coffee Brews. *J. Agric. Food Chem.* **2010**, *58*, 2958-2965.
13. Andueza, S.; Maeztu, L.; Dean, B.; de Pena, M.; Bello, J.; Cid, C. Influence of water pressure on the final quality of arabica espresso coffee. Application of multivariate analysis. *J. Agric. Food Chem.* **2002**, *50*, 7426-7431.

14. Andueza, S.; Maeztu, L.; Pascual, L.; Ibanez, C.; de Peña, M.; Cid, C. Influence of extraction temperature on the final quality of espresso coffee. *J. Sci. Food Agric.* **2003**, *83*, 240-248.
15. Andueza, S.; Vila, M.; de Peña, M.; Cid, C. Influence of coffee/water ratio on the final quality of espresso coffee. *J. Sci. Food Agric.* **2007**, *87*, 586-592.
16. Andueza, S.; De Peña, M.; Cid, C. Chemical and sensorial characteristics of espresso coffee as affected by grinding and torrefacto roast. *J. Agric. Food Chem.* **2003**, *51*, 7034-7039.
17. Cruz, R.; Cardoso, M.M.; Fernandes, L.; Oliveira, M.; Mendes, E.; Baptista, P.; Morais, S.; Casal, S. Espresso coffee residues: a valuable source of unextracted compounds. *J. Agric. Food Chem.* **2012**, *60*, 7777-7784.
18. Zuorro, A.; Lavecchia, R. Spent coffee grounds as a valuable source of phenolic compounds and bioenergy. *J. Cleaner Prod.* **2012**, *34*, 49-56.
19. Bravo, J.; Monente, C.; Juániz, I.; De Peña, M.P.; Cid, C. Influence of extraction process on antioxidant capacity of spent coffee. *Food Res. Int.* **2011**, DOI:10.1016/j.foodres.2011.04.026.
20. Trugo, L.C.; Macrae, R. A study of the effect of roasting on the chlorogenic acid composition of coffee using HPLC. *Food Chem.* **1984**, *15*, 219-227.
21. Farah, A.; De Paulis, T.; Trugo, L.; Martin, P. Effect of roasting on the formation of chlorogenic acid lactones in coffee. *J. Agric. Food Chem.* **2005**, *53*, 1505-1513.
22. Bicchi, C.P.; Binello, A.E.; Pellegrino, G.M.; Vanni, A.C. Characterization of Green and Roasted Coffees through the Chlorogenic Acid Fraction by HPLC-UV and Principal Component Analysis. *J. Agric. Food Chem.* **1995**, *43*, 1549-1555.
23. Meydav, S.; Saguy, I.; Kopelman, I.J. Browning determination in citrus products. *J. Agric. Food Chem.* **1977**, *25*, 602-604.
24. Singleton, V.; Rossi, J. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enol. Vitic.* **1965**, *16*, 144-158.
25. Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M.; Rice-Evans, C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biol. Med.* **1999**, *26*, 1231-1237.
26. Brand-Williams, W.; Cuvelier, M.E.; Berset, C. Use of a free radical method to evaluate antioxidant activity. *LWT -- Food Sci. Technol.* **1995**, *28*, 25-30.
27. Roesch, D.; Bergmann, M.; Knorr, D.; Kroh, L.W. Structure–Antioxidant Efficiency Relationships of Phenolic Compounds and Their Contribution to the Antioxidant Activity of Sea Buckthorn Juice. *J. Agric. Food Chem.* **2003**, *51*, 4233-4239.

28. Cämmerer, B.; Kroh, L. Antioxidant activity of coffee brews. *Eur. Food Res. Technol.* **2006**, *223*, 469-474.
29. Clifford, M.N. Chlorogenic acids. In *Coffee. Vol. 1. Chemistry*. Clarke, R.J. and Macrae, R., Eds.; Barking, Essex, UK; Elsevier Science Publishers, 1985; pp. 153-202.
30. Farah, A.; Donangelo, C.M. Phenolic compounds in coffee. *Braz. J. Plant Physiol.* **2006**, *18*, 23-36.
31. Vignoli, J.A.; Bassoli, D.G.; Benassi, M.T. Antioxidant activity, polyphenols, caffeine and melanoidins in soluble coffee: The influence of processing conditions and raw material. *Food Chem.* **2011**, *124*, 863-868.
32. Ludwig, I.A.; Sanchez, L.; Caemmerer, B.; Kroh, L.W.; De Peña, M.P.; Cid, C. Extraction of coffee antioxidants: Impact of brewing time and method. *Food Res. Int.* **2012**, *48*, 57-64.
33. Clifford, M.N. The nature of chlorogenic acids: Are they advantageous compounds in coffee?. In *Proceedings 17th International Scientific Colloquium on Coffee*. Nairobi, Kenia, **1997**, 79-91.
34. Perrone, D.; Donangelo, R.; Donangelo, C.M.; Farah, A. Modeling Weight Loss and Chlorogenic Acids Content in Coffee during Roasting. *J. Agric. Food Chem.* **2010**, *58*, 12238-12243.
35. Blumberg, S.; Frank, O.; Hofmann, T. Quantitative Studies on the Influence of the Bean Roasting Parameters and Hot Water Percolation on the Concentrations of Bitter Compounds in Coffee Brew. *J. Agric. Food Chem.* **2010**, *58*, 3720-3728.
36. Bekedam, E.K.; Schols, H.; Van Boekel, Martinus A J S.; Smit, G. Incorporation of chlorogenic acids in coffee brew melanoidins. *J. Agric. Food Chem.* **2008**, *56*, 2055-2063.
37. Nunes, F.; Coimbra, M. Role of hydroxycinnamates in coffee melanoidin formation. *Phytochem. Rev.* **2010**, *9*, 171-185.
38. Perrone, D.; Farah, A.; Donangelo, C.M. Influence of Coffee Roasting on the Incorporation of Phenolic Compounds into Melanoidins and Their Relationship with Antioxidant Activity of the Brew. *J. Agric. Food Chem.* **2012**, *60*, 4265-4275.
39. Leifa, F.; Pandey, A.; Soccol, C.R. Solid state cultivation -- an efficient method to use toxic agro-industrial residues. *J. Basic Microbiol.* **2000**, *40*, 187-197.
40. Belitz, H.; Grosch, W.; Schieberle, P. Coffee, Tea, Cocoa. In *Food Chemistry*, 4th ed.; Belitz, H., Grosch, W., Schieberle, P., Eds.; Springer-Verlag Berlin: Heidelberg, 2009; pp. 938-969.
41. López-Galilea, I.; Andueza, S.; Leonardo, I.d.; de Peña, M.; Cid, C. Influence of torrefacto roast on antioxidant and pro-oxidant activity of coffee. *Food Chem.* **2006**, *94*, 75-80.

42. Rufian-Henares, J.; Morales, F. Effect of in Vitro Enzymatic Digestion on Antioxidant Activity of Coffee Melanoidins and Fractions. *J. Agric. Food Chem.* **2007**, *55*, 10016-10021.
43. Delgado-Andrade, C.; Rufian-Henares, J.; Morales, F. Assessing the antioxidant activity of melanoidins from coffee brews by different antioxidant methods. *J. Agric. Food Chem.* **2005**, *53*, 7832-7836.
44. Dorea, J.; da Costa, T. Is coffee a functional food? *Br. J. Nutr.* **2005**, *93*, 773-782.
45. Crozier, A.; Jaganath, I.B.; Clifford, M.N. Dietary phenolics: chemistry, bioavailability and effects on health. *Nat. Prod. Rep.* **2009**, *26*, 1001-1043.
46. Murthy, P.S.; Madhava Naidu, M. Sustainable management of coffee industry by-products and value addition—A review. *Resour., Conserv. Recycl.* **2012**, *66*, 45-58.
47. Bekedam, E.K.; Schols, H.A.; Cämmerer, B.; Kroh, L.W.; van Boekel, Martinus A. J. S.; Smit, G. Electron Spin Resonance (ESR) Studies on the Formation of Roasting-Induced Antioxidative Structures in Coffee Brews at Different Degrees of Roast. *J. Agric. Food Chem.* **2008**, *56*, 4597-4604.
48. Delgado-Andrade, C.; Morales, F.J. Unraveling the Contribution of Melanoidins to the Antioxidant Activity of Coffee Brews. *J. Agric. Food Chem.* **2005**, *53*, 1403-1407.
49. Wang, H.; Qian, H.; Yao, W. Melanoidins produced by the Maillard reaction: Structure and biological activity. *Food Chem.* **2011**, *128*, 573-584.

FIGURE CAPTIONS

Figure 1. Antioxidant capacity percentage of Arabica (a) and Robusta (b) spent coffee obtained from different coffeemakers in comparison to their respective coffee brew (expressed as 100%).


F: Filter, E: Espresso, P: Plunger, M: Mocha.

Figure 2. Fremy's salt (a) and TEMPO (b) antioxidant capacity of spent coffee obtained from filter and espresso coffeemakers.

Figure 1. Antioxidant capacity percentage of Arabica (a) and Robusta (b) spent coffee obtained from different coffeemakers in comparison to their respective coffee brew (expressed as 100%).

F: Filter, E: Espresso, P: Plunger, M: Mocha.

a) Arabica



b) Robusta





Figure 2. Fremy's salt (a) and TEMPO (b) antioxidant capacity of spent coffee obtained from filter and espresso coffeemakers.

a) Fremy's salt



b) TEMPO




Table 1. Caffeoylquinic acids (mg/g) in spent coffee and coffee brews.

	Spent coffee				Coffee brews			
	Filter	Espresso	Plunger	Mocha	Filter	Espresso	Plunger	Mocha
3-CQA								
Arabica	1.64 ± 0.00 ^d	1.29 ± 0.03 ^c	1.10 ± 0.03 ^b	nd ^a	3.54 ± 0.00 ^{ab}	3.08 ± 0.07 ^a	3.91 ± 0.22 ^b	5.29 ± 0.29 ^c
Robusta	0.68 ± 0.03 ^b	0.83 ± 0.03 ^c	0.63 ± 0.02 ^b	nd ^a	2.26 ± 0.13 ^b	1.81 ± 0.09 ^a	2.03 ± 0.11 ^{ab}	3.43 ± 0.05 ^c
4-CQA								
Arabica	2.51 ± 0.02 ^d	2.03 ± 0.02 ^c	1.75 ± 0.10 ^b	nd ^a	5.50 ± 0.00 ^b	3.97 ± 0.05 ^a	6.01 ± 0.32 ^b	7.67 ± 0.41 ^c
Robusta	0.97 ± 0.04 ^b	1.16 ± 0.01 ^c	0.99 ± 0.05 ^b	nd ^a	2.93 ± 0.17 ^b	2.45 ± 0.02 ^a	3.01 ± 0.18 ^b	4.03 ± 0.02 ^c
5-CQA								
Arabica	3.59 ± 0.02 ^d	2.84 ± 0.04 ^c	2.48 ± 0.07 ^b	nd ^a	7.42 ± 0.01 ^a	6.89 ± 0.13 ^a	8.57 ± 0.43 ^b	10.72 ± 0.61 ^c
Robusta	1.26 ± 0.06 ^b	1.56 ± 0.05 ^c	1.18 ± 0.05 ^b	nd ^a	4.38 ± 0.21 ^b	3.70 ± 0.14 ^a	3.63 ± 0.20 ^a	6.25 ± 0.07 ^c
Total CQAs								
Arabica	7.74	6.16	5.33	nd	16.46	13.94	18.49	23.68
Robusta	2.91	3.55	2.80	nd	9.57	7.96	8.67	13.71
3,4-diCQA								
Arabica	2.53 ± 0.03 ^c	2.34 ± 0.02 ^b	2.46 ± 0.06 ^c	nd ^a	0.22 ± 0.00 ^a	0.47 ± 0.02 ^b	0.57 ± 0.03 ^c	0.89 ± 0.05 ^d
Robusta	1.49 ± 0.07 ^b	1.80 ± 0.01 ^c	1.99 ± 0.10 ^d	nd ^a	0.29 ± 0.01 ^a	0.29 ± 0.00 ^a	0.30 ± 0.02 ^a	0.52 ± 0.00 ^b
3,5-diCQA								
Arabica	1.17 ± 0.01 ^c	1.09 ± 0.02 ^b	1.26 ± 0.05 ^d	nd ^a	nd ^a	0.25 ± 0.01 ^c	0.22 ± 0.01 ^b	0.42 ± 0.02 ^d
Robusta	0.62 ± 0.03 ^b	0.74 ± 0.01 ^c	0.70 ± 0.04 ^c	nd ^a	0.18 ± 0.01 ^c	0.14 ± 0.00 ^a	0.16 ± 0.01 ^b	0.32 ± 0.00 ^d
4,5-diCQA								
Arabica	1.80 ± 0.03 ^c	1.46 ± 0.02 ^b	2.07 ± 0.10 ^d	nd ^a	nd ^a	0.35 ± 0.01 ^b	0.37 ± 0.02 ^b	0.63 ± 0.03 ^c
Robusta	1.20 ± 0.05 ^b	1.40 ± 0.03 ^c	1.41 ± 0.08 ^c	nd ^a	nd ^a	0.26 ± 0.00 ^b	0.26 ± 0.01 ^b	0.74 ± 0.00 ^c
Total diCQAs								
Arabica	5.50	4.89	5.79	nd	0.22	1.07	1.16	1.94
Robusta	3.31	3.94	4.10	nd	0.47	0.69	0.72	1.58
Total CQAs+diCQAs								
Arabica	13.24	11.05	11.12	nd	16.68	15.01	19.65	25.62
Robusta	6.22	7.49	6.90	nd	10.04	8.65	9.39	15.29

All values are shown as means ± SD (n=3). In each row, different superscripts indicate significant differences (p<0.05) among different spent coffee or coffee brews.

CQA, monocaffeoylquinic acids; diCQA, dicaffeoylquinic acids. nd, not detected.

Table 2. Caffeine (mg/g) and melanoidins (Abs 420nm) in spent coffee and coffee brews.

	Spent coffee				Coffee brews			
	Filter	Espresso	Plunger	Mocha	Filter	Espresso	Plunger	Mocha
Caffeine								
Arabica	5.20 ± 0.04 ^d	3.59 ± 0.00 ^b	3.76 ± 0.12 ^c	nd ^a	10.08 ± 0.04 ^a	10.07 ± 0.17 ^a	11.83 ± 0.65 ^b	15.64 ± 0.86 ^c
Robusta	7.53 ± 0.35 ^c	8.09 ± 0.05 ^c	5.73 ± 0.31 ^b	nd ^a	22.08 ± 0.06 ^c	17.73 ± 0.14 ^b	15.64 ± 0.84 ^a	22.68 ± 0.02 ^c
Melanoidins								
Arabica	0.165 ± 0.006 ^d	0.102 ± 0.001 ^b	0.114 ± 0.001 ^c	tr ^a	0.155 ± 0.003 ^a	0.315 ± 0.008 ^b	0.401 ± 0.009 ^c	0.803 ± 0.011 ^d
Robusta	0.145 ± 0.005 ^d	0.133 ± 0.001 ^c	0.118 ± 0.000 ^b	tr ^a	0.197 ± 0.001 ^a	0.418 ± 0.008 ^b	0.647 ± 0.006 ^c	0.847 ± 0.009 ^d

All values are shown as means ± SD (n=3). In each row, different superscripts indicate significant differences (p<0.05) among different spent coffee or coffee brews. nd, not detected. tr, traces.

Table 3. Antioxidant capacity of spent coffee and coffee brews.

	Spent coffee				Coffee brews			
	Filter	Espresso	Plunger	Mocha	Filter	Espresso	Plunger	Mocha
Folin Ciocalteu (mg GA/g)								
Arabica	24.60 ± 0.18 ^d	15.79 ± 0.08 ^c	13.87 ± 0.16 ^b	1.36 ± 0.03 ^a	26.70 ± 0.07 ^b	23.13 ± 0.23 ^a	33.29 ± 0.13 ^c	50.16 ± 0.07 ^d
Robusta	17.54 ± 0.26 ^c	19.12 ± 0.14 ^d	13.22 ± 0.20 ^b	1.88 ± 0.05 ^a	38.16 ± 1.31 ^b	30.55 ± 0.06 ^a	51.41 ± 1.02 ^c	60.92 ± 0.46 ^d
ABTS (µmol Trolox/g)								
Arabica	215.12 ± 2.18 ^d	131.83 ± 1.49 ^c	120.98 ± 1.81 ^b	8.09 ± 0.33 ^a	242.08 ± 3.54 ^b	206.12 ± 1.68 ^a	280.11 ± 7.01 ^c	456.33 ± 5.79 ^d
Robusta	167.24 ± 5.78 ^c	177.63 ± 1.23 ^d	118.23 ± 3.05 ^b	22.17 ± 0.30 ^a	310.93 ± 2.27 ^b	299.86 ± 2.69 ^a	493.97 ± 1.27 ^c	622.45 ± 6.16 ^d
DPPH (µmol Trolox/g)								
Arabica	112.06 ± 2.21 ^d	74.57 ± 2.99 ^c	60.96 ± 1.75 ^b	2.56 ± 0.18 ^a	109.49 ± 2.80 ^b	87.13 ± 0.40 ^a	145.94 ± 3.66 ^c	208.18 ± 9.39 ^d
Robusta	74.09 ± 1.00 ^c	83.59 ± 2.39 ^d	53.99 ± 0.25 ^b	5.18 ± 0.06 ^a	128.18 ± 5.20 ^a	121.72 ± 4.69 ^a	231.06 ± 8.64 ^b	243.46 ± 0.75 ^b

All values are shown as means ± SD (n=3). In each row, different superscripts indicate significant differences ($p < 0.05$) among different spent coffee or coffee brews.